
(Revised version dated 16 February 2020)

Guidelines for medical practice, diagnosis, treatment and prevention of Healthcare-associated infection in response to patients with COVID-19 infection

For physicians and healthcare personnel



Negative for COVID-19 (1 laboratory)

- 1) Consider providing treatment as appropriate
- Treat as OPD case if no indication for hospitalization and personal health practices recommended, repeat the testing if not if improve
- If the conditions not improved within 48 hours, consider repeating testing for SAR-CoV2 at the reference laboratory

Positive for COVID-19

- Patient is admitted in single isolation room or cohort ward (designated specifically for confirmed case of COVID-19 infection) with at least 1-meter distance between the beds
- Admitted in AIIR for severe case, or in case aerosolgenerating procedure will need to be conducted.
- 3) Supportive treatment provided
- 4) Consult SMEs
- 5) To discharge the patient, considering the symptom and having negative laboratory result from 2 different laboratories; local laboratory and reference laboratory and repeat the testing in 48 hours at the local laboratory after the first testing

Guidelines for medical practice, diagnosis, treatment and prevention of healthcare-associated infection in response to patients with COVID-19 infection for physician and healthcare personnel by Department of Medical Services, Ministry of Public Health and a panel of subject matter experts from local medical schools.