

**COVID-19 Patient Under Investigation (PUI) definition: 29 January 2021**

Signs and Symptoms	Risk factors	Quarantine measures in case of negative test result
<p><b>Scenario 1: <u>Surveillance at Points of Entry</u></b>                      Patient with body temperature <math>\geq 37.3</math> °C or any of the following respiratory symptoms such as cough, runny nose, sore throat, anosmia, ageusia, tachypnea, or shortness of breath, or difficulty Breathing</p>	Having <u>history</u> of travel to or from foreign countries from all flights and points of entry	Follow quarantine measures
<p><b>Scenario 2: <u>Surveillance in PUIs/patients</u></b></p> <p><b>2.1</b> Symptomatic PUIs with any of the following symptoms: history of fever/body temperature <math>\geq 37.5</math> °C, cough, runny nose, sore throat, anosmia, ageusia, tachypnea, or shortness of breath, or difficulty breathing</p>	1) Any history within 14 days prior to symptom onset as follows: 1.1) Travel history to/ from or reside in areas <u>with ongoing local transmission in the past month</u> 1.2) History of contact with a confirmed COVID-19 case 1.3) Travel history to crowded places in the community or places of gathering <u>where confirmed cases have been reported in the past month</u> e.g., flea markets, malls, hospitals, public transportation 1.4) Work in a quarantine facility 2) Suspected of having COVID-19 by attending physician	-Follow quarantine measures for 1.1) in every case 1.2) and 1.3) in high-risk contacts  - For other cases, follow CPG of disease diagnosed
<p><b>2.2</b> Patients with pneumonia</p>	Any of the following characteristics: 1) Severe symptoms requiring intubation or death 2) Etiology unclear or cannot be identified within 48 hrs. 3) Suspected of having COVID-19 by attending physician	Follow CPG of disease diagnosed
<p><b>Scenario 3: <u>Surveillance in health care workers</u></b>                      with any of the following symptoms: history of fever/body temperature <math>\geq 37.5</math> °C, cough, runny nose, sore throat, anosmia, ageusia, tachypnea, or shortness of breath, or difficulty breathing</p>	Working in health care facilities e.g. hospitals, clinics, health promotion hospitals, laboratories, drug stores, members of investigation teams or personnel working in quarantine facilities (test as determined appropriate)	Follow CPG of disease diagnosed
<p><b>Scenario 4: <u>Surveillance in clusters of patients with respiratory infection</u></b>                      A cluster of <math>\geq 5</math> patients with symptoms of respiratory infection in the same specific area</p>	Clusters in the same area, in the same week with epidemiological linkage.	Follow CPG of disease diagnosed